VOL. 476 JOURNAL OF CHROMATOGRAPHY AUGUST 4, 1989

CONTENTS

8TH INTERNATIONAL SYMPOSIUM ON HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY OF PROTEINS, PEPTIDES AND POLYNUCLEOTIDES, COPENHAGEN, OCTOBER 31–NOVEMBER 2, 1988

Foreword
by M. T. W. Hearn 1

Fast separation of biological macromolecules on non-porous, microparticulate columns
by G. P. Rozing and H. Goetz (Waldbronn, F.R.G.) 3

New porous organic microspheres for high-performance liquid chromatography
by R. W. Stout, H. J. Leibu, A. T. Rousak and R. C. Wright (Wilmington, DE, U.S.A.) . 21

Evaluation of advanced silica packings for the separation of biopolymers by high-performance liquid chromatography. VI. Design, chromatographic performance and application of non-porous silica-based anion exchangers
by G. Jilge, K. K. Unger, U. Esser, H.-J. Schäfer and G. Rathgeber (Mainz, F.R.G.) and W. Müller (Darmstadt, F.R.G.) 37

Polymer support synthesis. XV. Behaviour of non-porous surface-coated silica gel microbeads in oligonucleotide synthesis
by H. Seliger, U. Kotschi, C. Scharpf and R. Martin (Ulm, F.R.G.), F. Eisenbeiss and J. N. Kinkel (Darmstadt, F.R.G.) and K. K. Unger (Mainz, F.R.G.) 49

Blotting of proteins onto Immobilon membranes. *In situ* characterization and comparison with high-performance liquid chromatography
by T. Choli, U. Kapp and B. Wittmann-Liebold (Berlin, F.R.G.) 59

Automated enantioseparation of amino acids by derivatization with *o*-phthaldialdehyde and N-acylated cysteines
by H. Brückner and R. Wittner (Stuttgart, F.R.G.) and H. Godel (Waldbronn, F.R.G.) . 73

Liquid chromatographic analysis of amino and imino acids in protein hydrolysates by post-column derivatization with *o*-phthalaldehyde and 3-mercaptopropionic acid
by A. Fiorino, G. Frigo and E. Cucchetti (Corsico, Italy) 83

High-performance liquid chromatography for cyclosporin measurement: comparison with radioimmunoassay
by M. Plebani, M. Masiero, C. D. Paleari, L. Sciacovelli, D. Faggian and A. Burlina (Padova, Italy) 93

Isolation of recombinant partial *gag* gene product p18 (HIV-1_{Bru}) from *Escherichia coli*
by H. V. J. Kolbe, F. Jaeger, P. Lepage, C. Roitsch, G. Lacaud, M.-P. Kieny, J. Sabatie, S. W. Brown and J.-P. Lecocq (Strasbourg, France) and M. Girard (Marnes-la-Coquette, France) 99

High-performance liquid chromatography of amino acids, peptides and proteins. XC. Investigations into the relationship between structure and reversed-phase high-performance liquid chromatography retention behaviour of peptides related to human growth hormone
by A.W. Purcell, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 113

High-performance liquid chromatography of amino acids, peptides and proteins. XCI. The influence of temperature on the chromatographic behaviour of peptides related to human growth hormone
by A. W. Purcell, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 125

Retention behaviour of paracelsin peptides on reversed-phase silicas with varying *n*-alkyl chain length and ligand density
by K. D. Lork and K. K. Unger (Mainz, F.R.G.), H. Brückner (Stuttgart, F.R.G.) and M. T. W. Hearn (Clayton, Australia) 135

Sodium dodecyl sulphate–protein complexes. Changes in size of shape below the critical micelle concentration, as monitored by high-performance agarose gel chromatography
by E. Mascher and P. Lundahl (Uppsala, Sweden) 147

Protein conformation changes as the result of binding to reversed-phase chromatography column materials
by A. F. Drake, M. A. Fung and C. F. Simpson (London, U.K.) 159

Stereochemical recognition of enantiomeric and diastereomeric dipeptides by high-performance liquid chromatography on a chiral stationary phase based upon immobilized α-chymotrypsin
by P. Jadaud and I. W. Wainer (Memphis, TN, U.S.A.) 165

Efficient endotoxin removal with a new sanitizable affinity column: Affi-Prep Polymyxin
by K. W. Talmadge and C. J. Siebert (Richmond, CA, U.S.A.) 175

Isolation of a specific membrane protein by immunoaffinity chromatography with biotinylated antibodies immobilized on avidin-coated glass beads
by J. V. Babashak (Vineland, NJ, U.S.A.) and T. M. Phillips (Washington, DC, U.S.A.) . 187

Coated silica supports for high-performance affinity chromatography of proteins
by F. L. Zhou, D. Muller, X. Santarelli and J. Jozefonvicz (Villetaneuse, France) 195

High-performance liquid chromatography of amino acids, peptides and proteins. XCII. Thermodynamic and kinetic investigations on rigid and soft affinity gels with varying particle and pore sizes
by F. B. Anspach, A. Johnston and H.-J. Wirth (Clayton, Australia), K. K. Unger (Mainz, F.R.G.) and M. T. W. Hearn (Clayton, Australia) 205

Affinity chromatography of recombinant *Rhizomucor miehei* aspartic proteinase on Si-300 bacitracin
by S. B. Mortensen, L. Thim, T. Christensen, H. Woeldike, E. Boel, K. Hjortshoej and M. T. Hansen (Bagsvaerd, Denmark) 227

Immunological-chromatographic analysis of lysozyme variants
by L. J. Janis, A. Grott and F. E. Regnier (West Lafayette, IN, U.S.A.) and S. J. Smith-Gill (Bethesda, MD, U.S.A.) 235

Isolation of recombinant hirudin by preparative high-performance liquid chromatography
by R. Bischoff, D. Clesse, O. Whitechurch, P. Lepage and C. Roitsch (Strasbourg, France) 245

Comparison of protein A, protein G and copolymerized hydroxyapatite for the purification of human monoclonal antibodies
by A. Jungbauer, C. Tauer, M. Reiter, M. Purtscher, E. Wenisch, F. Steindl, A. Buchacher and H. Katinger (Vienna, Austria) 257

High-performance liquid chromatography of amino acids, peptides and proteins. XCIII. Comparison of methods for the purification of mouse monoclonal immunoglobulin M autoantibodies
by G. Coppola, J. Underwood, G. Cartwright and M. T. W. Hearn (Clayton, Australia) . 269

Improved separation of human pepsins from gastric juice by high-performance ion-exchange chromatography
by K. Peek, N. B. Roberts and W. H. Taylor (Liverpool, U.K.) 291

Separation and purification of component proteins of the cytochrome P-450-dependent microsomal monooxygenase system by high-performance liquid chromatography
by H. Taniguchi and W. Pyerin (Heidelberg, F.R.G.) 299

Crown ethers as ligands for high-performance liquid chromatography of proteins and nucleic acids
by D. Josić, W. Reutter and J. Reusch (Berlin, F.R.G.) 309

Comparison of non-ionic detergents for extraction and ion-exchange high-performance liquid chromatography of Sendai virus integral membrane proteins
by J. van Ede, J. R. J. Nijmeijer and S. Welling-Wester (Groningen, The Netherlands), C. Örvell (Stockholm, Sweden) and G. W. Welling (Groningen, The Netherlands) 319

Properties, in theory and practice, of novel gel filtration media for standard liquid chromatography
by L. Hagel, H. Lunström, T. Andersson and H. Lindblom (Uppsala, Sweden) 329

Peptide mapping and internal sequencing of proteins electroblotted from two-dimensional gels onto polyvinylidene difluoride membranes. A chromatographic procedure for separating proteins from detergents
by R. J. Simpson, L. D. Ward, G. E. Reid, M. P. Batterham and R. L. Moritz (Melbourne, Australia) 345

Correlation of protein retention times in reversed-phase chromatography with polypeptide chain length and hydrophobicity
by C. T. Mant, N. E. Zhou and R. S. Hodges (Edmonton, Canada) 363

Strong cation-exchange high-performance liquid chromatography of peptides. Effect of non-specific hydrophobic interactions and linearization of peptide retention behaviour
by T. W. L. Burke, C. T. Mant, J. A. Black and R. S. Hodges (Edmonton, Canada) . . . 377

High-performance liquid chromatography of amino acids, peptides and proteins. LXXXIX. The influence of different displacer salts on the retention properties of proteins separated by gradient anion-exchange chromatography
by A. N. Hodder, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 391

Improved separation speed and efficiency for proteins, nucleic acids and viruses in asymmetrical flow field flow fractionation
by A. Litzén and K.-G. Wahlund (Uppsala, Sweden) 413

Ester and related derivatives of ring N-pentafluorobenzylated 5-hydroxymethyluracil. Hydrolytic stability, mass spectral properties, and trace detection by gas chromatography–electron-capture detection, gas chromatography–electron-capture negative ion mass spectrometry, and moving-belt liquid chromatography–electron-capture negative ion mass spectrometry
by G. M. Kresbach, M. Itani, M. Saha, E. J. Rogers, P. Vouros and R. W. Giese (Boston, MA, U.S.A.) 423

Purification of synthetic oligonucleotides on a weak ion-exchange column
by J. Liautard, C. Ferraz, J. Sri Widada, J. P. Capony and J. P. Liautard (Montpellier, France) 439

High-performance liquid chromatography of amino acids, peptides, proteins and polynucleotides. XCIV. Solid-phase hybridization of complementary oligonucleotides
by M. Guthridge, J. Bertolini and M. T. W. Hearn (Clayton, Australia) 445

Separation of two molecular forms of human estrogen receptor by hydrophobic interaction chromatography. Gradient optimization and tissue comparison
by S. M. Hyder and J. L. Wittliff (Louisville, KY, U.S.A.) 455

High-performance liquid chromatofocusing and column affinity chromatography of *in vitro* ^{14}C-glycated human serum albumin. Demonstration of a glycation-induced anionic heterogeneity
by P. Vidal, T. Deckert, B. Hansen and B. S. Welinder (Gentofte, Denmark) 467

Comparison of ion-exchange high-performance liquid chromatography columns for purification of Sendai virus integral membrane proteins
by S. Welling-Wester, R. M. Haring and H. Laurens (Groningen, The Netherlands), C. Örvell (Stockholm, Sweden) and G. W. Welling (Groningen, The Netherlands) 477

High-performance liquid chromatographic assay for nicotinamide–adenine dinucleotide kinase
by M. Pace, P. L. Mauri, C. Gardana and P. G. Pietta (Milan, Italy) 487

Heterogeneity of human pepsin 1, as shown by high-performance ion-exchange chromatography
by K. Peek, N. B. Roberts and W. H. Taylor (Liverpool, U.K.) 491

Reversed-phase liquid chromatography of proteins with strong acids
by G. Thévenon and F.E. Regnier (West Lafayette, IN, U.S.A.) 499

Author Index 513